

Lu et al. reviewed possible entry pathways of the oral-gut microbiome axis, and the influence of oral microbes, on intestinal microbiota and the immune system. With the popularity of oral-gut microbes research, studying the influence of oral microbes on the intestine opens up a new perspective into preventing and managing intestinal diseases.

In another review article, Gershater et al. reviewed the characteristic of the oral microbiome of patients with cleft lip and palate (CL/P). The review shed light on the significant microbiota differences between CL/P patients and healthy individuals in different anatomic regions, including the teeth inside and adjacent to the cleft, oral cavity, nasal cavity, pharynx, and ear, as well as bodily fluids, secretions, and excretions. The study would be conducive to the advance of CL/P-specific microbiota treatment. Interestingly, the feature of a given species of CL/P patients and healthy individuals was not consistent from different sampling regions, suggesting that it must be done to sufficiently explore the non-typical bacterial and fungal species of CL/P patients in the future.

The oral microbiota may be a potential target for treating cardiovascular diseases (Tonelli et al., 2023). Hernandez-Ruiz et al. evaluated oral microbiota diversity and circulating inflammatory profile in ST-segment elevation myocardial infarction (STEMI) patients stratified according to an inflammation-based risk scoring system and defined a non-causal link inferred between the cardiovascular risk of STEMI patients, and its relationship with the exacerbation of the inflammation. The study demonstrated that Bacteroidetes phylum was most abundant in STEMI patients, and Prevotella was the most abundant genus, with a higher proportion in periodontitis patients. Prevotella genus was found to correlate positively and significantly with elevated IL-6 concentration. The study emphasized the relationship between oral microbiome and circulatory system disease.

The balance of the microbial community is vital to infant health and development. Hou et al. developed a novel decontamination method for the new ultra-sensitive metagenomic sequencing method named 2bRAD-M and identified 11 and 8 potential biomarkers for neonatal bacterial meningitis in blood and cerebrospinal fluid (CSF). They further revealed 16 and 35 microbial species that were highly associated with the physiological indicators in blood and CSF, respectively. The study offered new microbiological markers to assist in the diagnosis of neonatal bacterial meningitis.

Also, Li et al. provided the future direction to apply the oral microbiome to improve human health. They depicted a synthetic and exhaustive landscape of focus and forefront in oral microbiome studies. The analysis suggested that the homeostatic balance of the oral microbiome, advanced microbial sequencing technology, connections with gut microbiota, and tumorigenesis have become hotspots in the aspect of the oral microbiome. Their results shed

light on the increasing importance of the oral microbiome in the perspective of systemic diseases and states, more diverse fields such as immunity and cancer. Moreover, the study encouraged to prioritize fostering cooperation between institutions and researchers, both nationally and internationally.

Conclusion

Evidence linking the oral microbiota to systemic diseases and overall health continues to accumulate. The physiological and pathological interaction between oral microbiome and microbiomes of other body sites has been suggested. A better understanding and effective management of oral microbiome-related diseases call for a holistic view of interactions among different host microbiota which requires systemic and multifactorial approaches.

Author contributions

TX: Writing original draft. NC: Supervision, Writing review & editing. XH: Supervision, Writing review & editing. FC: Supervision, Writing review & editing.

Acknowledgments

We thank all the contributors of this Research Topic.

Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

The authors declared that they were an editorial board member of Frontiers, at the time of submission. This had no impact on the peer review process and the final decision.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

References

- Gao, Lu, Xu, T., Huang, G., Jiang, S., Gu, Y., and Chen, F. (2018). Oral microbiomes: more and more importance in oral cavity and whole body. *Protein Cell* 9 (5), 488–500. doi: 10.1007/s13238-018-0548-1
- Hou, K., Wu, Z.-X., Chen, X.-Y., Wang, J.-Q., Zhang, D., Xiao, C., et al. (2022). Microbiota in health and diseases. *Signal Transduct. Target. Ther.* 7 (1), 135–135. doi: 10.1038/s41392-022-00974-4
- Lamont, R. J., Koo, H., and Hajishengallis, G. (2018). The oral microbiota: dynamic communities and host interactions. *Nat. Rev. Microbiol.* 16 (12), 745–759. doi: 10.1038/s41579-018-0089-x
- Tonelli, A., Lumngwena, E. N., and Ntusi, N. A. B. (2023). The oral microbiome in the pathophysiology of cardiovascular disease. *Nat. Rev. Cardiol.* 20 (6), 386–403. doi: 10.1038/s41569-022-00825-3
- Tuganbaev, T., Yoshida, K., and Honda, K. (2022). The effects of oral microbiota on health. *Science* 376 (6596), 934–936. doi: 10.1126/science.abn1890